

Prevalence, species distribution, and risk factors of fungal colonization and infection in patients at a burn intensive care unit in Vietnam

Nguyen Van Bang^{1#}, Nguyen Thanh Xuan^{2#}, Dinh Xuan Quang³, Cao Ba Loi³, Nguyen Thai Ngoc Minh⁴, Nguyen Nhu Lam⁴, Do Ngoc Anh⁵, Truong Thi Thu Hien⁶, Hoang Xuan Su⁷, Le Tran-Anh^{5*}

¹ Department of Hamatology, Toxicology, Radiation, and Occupational Diseases, Military Hospital 103, Vietnam Military Medical University, Ha Dong, Ha Noi, Vietnam

² Department of Medical Education, Military Hospital 103, Vietnam Military Medical University, Ha Dong, Ha Noi, Vietnam

³ Department of Scientific and Training Management, National Institute of Malaria, Parasitology, and Entomology, Nam Tu Liem, Ha Noi, Vietnam

⁴ Intensive Care Unit, National Hospital of Burn, Vietnam Military Medical University, Ha Dong, Ha Noi, Vietnam

⁵ Department of Parasitology, Vietnam Military Medical University, Ha Dong, Ha Noi, Vietnam

⁶ Department of Microbiology, National Hospital of Burn, Vietnam Military Medical University, Ha Dong, Ha Noi, Vietnam

⁷ Department of Microbiology and Pathogens, Institute of Biomedicine and Pharmacy, National Hospital of Burn, Vietnam Military Medical University, Ha Dong, Ha Noi, Vietnam

These authors contributed equally to this work and acted as joint first authors.

Article Info	ABSTRACT
Article type: Original article	Background and Purpose: Burn patients are at a higher risk of infections caused by different organisms. This study aimed to address the prevalence, causative species, and factors related to fungal colonization or infection in patients with acute severe injuries admitted to the intensive care unit (ICU) of a burn hospital in northern Vietnam.
Article History: Received: 19 August 2020 Revised: 29 September 2020 Accepted: 21 October 2020	Materials and Methods: This prospective study was conducted on 400 patients in a burn ICU between 2017 and 2019. Clinical samples were weekly collected and screened for fungi, and relevant clinical information was obtained from medical records.
* Corresponding author: Le Tran-Anh Department of Parasitology, Vietnam Military Medical University, Ha Dong, Ha Noi, Vietnam. Email: anh_lt@vmmu.edu.vn	Results: According to the results, 90% of the patients were colonized with fungi. Out of this group, 12.75% of the cases had invasive fungal infection (IFI). Eleven yeasts and six mold species were isolated from the patients, with the most common species being <i>Candida tropicalis</i> (45.56%) and <i>C. albicans</i> (41.94%). Among the eleven species causing fungal wound infection (FWI), the most common agents were <i>Candida</i> (66.7% of FWI patients) and <i>Aspergillus</i> (38.5%) species. Three <i>Candida</i> species isolated from blood were <i>C. tropicalis</i> (66.7%), <i>C. albicans</i> (20.0%), and <i>C. parapsilosis</i> (14.3%). No factors were found to expose the patients to a higher risk of fungal colonization. However, hyperglycemia, prolonged ICU stay, and heavy <i>Candida</i> species colonization were found to be independently predictive of IFI.
	Conclusion: Burn patients are at the risk of fungal infection with <i>Candida</i> species (especially <i>C. tropicalis</i>) and <i>Aspergillus</i> as the most frequently responsible agents. Continuous surveillance of fungi and appropriate management of pathophysiological consequences are essential to prevent fungal infection in burn patients.
	Keywords: Fungal colonization, Fungal infection, Burn patients, Intensive care unit, Vietnam

➤ How to cite this paper

Van Bang N, Thanh Xuan N, Xuan Quang D, Ba Loi C, Thai Ngoc Minh N, Nhu Lam N, Ngoc Anh D, Thi Thu Hien T, Xuan Su H, Tran-Anh L. Prevalence, species distribution, and risk factors of fungal colonization and infection in patients at a burn intensive care unit in Vietnam. *Curr Med Mycol.* 2020; 6(3): 42-49. DOI: [10.18502/cmm.6.3.4664](https://doi.org/10.18502/cmm.6.3.4664)

Introduction

Burn injuries are among the most common and detrimental types of all injuries and account for 180,000 deaths annually [1]. The leading cause of mortality among burned patients is infection (75%) [2]. Burn wounds provide an ideal port of entry for organisms and induce substantial immune dysfunction [3]. Accordingly, burn patients are at a higher risk of developing infections, compared with other hospitalized patients [4]. The

most common agents of infection among burn patients are bacteria and fungi [2, 5]. Due to the better care of burn patients and availability of effective antibiotics, there is a decreasing trend in infection due to drug-sensitive bacteria. However, the number of infections caused by multi-drug resistant bacteria or fungi is on an increasing trend [5, 6].

Fungal infections usually occur after a period of colonization with fungal strains either from hospital

environment or patients [7-10]. The effective prevention and treatment approaches of fungal infection should be based on updated and local epidemiological data because the prevalence, species distribution, and factors related to colonization or infection vary from time to time and place to place [11].

Vietnam is a developing country; regarding this, people living there are at a higher risk of burn injuries, compared to those living in developed countries [12]. However, no data on fungal colonization and infection among burn patients in Vietnam have been published. Regarding this, the aim of the present study was to address the prevalence, causative species, and factors related to fungal colonization or infection in patients with acute severe injuries admitted to the intensive care unit (ICU) of a burn hospital in northern Vietnam.

Materials and Methods

Ethics statement

The study was approved by the Ethics Committee of Vietnam National Institute of Malariology, Parasitology and Entomology (Decision number 1172/QD-VSR, 28th October 2016). Written informed consent was obtained for participation in the study. In cases where patients were not able to give consent themselves, the consent was obtained from their legal representatives.

Patients and data collection

The current investigation was conducted in the National Hospital of Burn (NHB), a teaching hospital of Vietnam Military Medical University (VMMU) during January 2017 and December 2019. Agent isolation and species identification were performed in the laboratories of the Department of Microbiology, NHB and Department of Parasitology, VMMU. All patients in the ICU of NHB during 2017-2019 who agreed to participate in this prospective study were enrolled in the research. The inclusion criteria were admission within 7 days of injury and presence of a thermal burn.

Burn injuries were managed at the NHB according to standard care for severe burn patients, such as aggressive resuscitation, use of silver sulfadiazine cream, wet wound dressing, surgical excision, and empirical antibiotic therapy. The data were obtained from inpatient charts and electronic medical records using a standardized case report form. Data collection was performed by senior clinicians who were blind to the study purpose and design. The Wallace rule was used to estimate the percentage of the total body surface area burned (% TBSA) and full-thickness surface area of patients [13]. The acute physiology and chronic health evaluation (APACHE) was used upon ICU admission to assess the injury severity for each patient [14].

Fungal surveillance

Non-sterile samples (i.e., mouth swabs, rectum swabs/feces, wound swabs, urine, and tracheal

aspirates) were collected weekly to detect the presence of microorganisms as part of the hospital standard of care. Wound biopsy and blood were taken in cases with clinically suspected fungal wound infection (FWI) (presence of pus, increased exudation, leathery appearance, change in wound color, inflammation, or graft lysis) or fungemia [2, 15]. The non-sterile samples were taken by two ready-made sterile cotton swabs. Blood was collected from a clean, unburnt site following routine protocols. Furthermore, biopsy (a 0.5×0.5-cm full-thickness sample) was taken from the most apparently infected site of the wound up to the healthy layer and then transferred to the laboratory in normal saline and 10% formaldehyde solution.

Fluids and secretions were microscopically examined after being processed by 10% potassium hydroxide. The biopsy samples fixed in 10% formaldehyde solution were sent to the pathology laboratory for hematoxylin and eosin (H&E) and periodic acid shift (PAS) staining, followed by an examination by pathologists. Non-sterile and tissue samples were cultured onto Sabouraud dextrose agar (SDA, BioMerieux, France) and slants containing gentamicin (0.02 mg/mL) to test the fungal growth. Blood samples were placed in the Bactec (BD Diagnostic Systems, Sparks, MD) system and daily checked for fungal growth.

The identification of the yeast isolates was accomplished by comparing the features of colonies on Brilliance Candida Agar (Oxoid) using the Vitek2 Compact (Biomérieux, France) following the manufacturer's instruction. Molds were identified based on morphological characteristics with the help of commonly used keys. The strains isolated from the sterile samples were further reconfirmed by sequencing the internal transcribed spacer (ITS) regions, and the obtained nucleotides were compared to database sequences in GenBank [16].

Definitions

Fungal colonization (FC) was defined as the isolation of fungi from a non-sterile body site, while invasive fungal infection (IFI) was referred to the presence of fungal elements in sterile sites (wound biopsy or blood) [17]. Colonization index (CI) was defined as the ratio of the number of distinct non-sterile samples colonized by *Candida* to the total number of body sites cultured; accordingly, the patients with a CI of ≥ 0.5 were considered heavily colonized [18].

The adult (aged 16-65 years) who had the TBSA of $\geq 50\%$ or FTBS of $\geq 10\%$ were considered severely burned. The patients aged < 16 or > 65 years with a TBSA of $\geq 30\%$ or an FTBS of $\geq 5\%$ were defined as severe burn patients. All patients with respiratory burns or burns associated with multiple injuries were also considered to have a severe injury.

The diagnosis of renal failure was based on variations in creatinine and urine output according to the risk, injury, failure, loss of function, and end-stage

renal disease criteria [19]. In addition, the patients with a fasting glucose level of > 126 mg/dl were considered to have hyperglycemia. The diagnosis of severe infection was performed based on the international guidelines for the treatment of sepsis and septic shock [20]. The patients who stayed in ICU for more than 14 days were regarded as having prolonged ICU stay [21].

Statistical analysis

The data were analyzed in SPSS software, version 16.0. Continuous variables were reported by means and standard deviation and compared by Student's t-test. Categorical variables were expressed as case number (n) and percentages and compared by the Chi-square test or Fisher's exact test as appropriate. Univariate analysis (linear regression) was used to investigate the possible correlations of independent variables with dependent ones (FC or IFI). Factors with a p-value of < 0.05 in univariate analysis were included in the model of the multivariate analysis. Odds ratios (OR) and corresponding 95% confidence intervals (95%CI) were calculated. Significance was set at a p-value of 0.05 (two-tailed).

Results

During the study time, there were 400 patients with a mean age of 29.7 years (range: 1-94) included in the research. The two most affected age groups were 1 ~ 10 (25%) and 31 ~ 40 (23.0%) years. Based on the obtained male: female ratio (3.5/1), males were found to be more affected than females. A total of 69 patients had inhalation injury, and 78 patients passed away during their ICU stay (Table 1). All infected patients were also colonized with fungi; accordingly, 360 (90%) patients were diagnosed with FC. Among 51 (12.25%) patients diagnosed with IFI, there were 3 patients with isolations from both blood culture and

tissue biopsy (Table 2).

The frequency of fungal species isolated from burned patients is listed in Table 3. Among the 11 identified yeast species, *C. tropicalis*, *C. albicans*, and *C. parapsilosis* were the most prevalent ones, accounting for 94.6% of all isolates. Some representative ITS2 sequences of the isolates were submitted to Genbank (*C. tropicalis* MN067757, *C. albicans* MT193530, *C. parapsilosis* MN067761, *C. glabrata* MN067758, *C. duobushaemulonii* MN174034, *Kodamaea ohmeri* MN067739, *A. oryzae* MH934917, *A. fumigatus* MH932411, *A. flavus* MN173209, *A. chevalieri* MN174037, *A. nomius* MN174036, and *F. solani* MN066126).

Table 1. Demographic characteristics of the patients

Characteristics	n	%	
Age groups (years)	1-10	100	25.00
	11-20	34	8.50
	21-30	62	15.50
	31-40	92	23.00
	41-50	52	13.00
	51-60	31	7.75
61-94	29	7.25	
Mean of age ($\bar{x} \pm SD$)	29.74 ± 20.86		
Gender	Male	311	77.75
	Female	89	22.25
Severe infection	327	81.75	
Severe burn	310	77.50	
Hyperglycemia	171	42.75	
Renal failure	81	20.25	
Inhalation injury	69	17.25	
APACHE>20	31	7.75	
Urinary catheter	393	98.25	
Prolonged ICU stay	369	92.25	
Parenteral nutrition	300	75.00	
Central venous catheters	276	69.00	
Immunosuppressive therapy	210	52.50	
Mechanical ventilation	124	31.00	
Hemodialysis	59	14.75	
Mortality	78	19.50	

APACHE: acute physiology and chronic health evaluation

Table 2. Prevalence of fungal colonization/infection (n=400)

Fungal colonization/infection			n	Percentage (%)
Yes	FC	Yeasts	22	5.50
		Molds	11	2.75
	Invasive fungal infection	Yeasts and molds	3	0.75
		Fungemia	12	3.00
		Fungemia+FWI	3	0.75
	No		40	10.00

FC: fungal colonization, FWI: fungal wound infection

Table 3. Distribution of the isolated yeast species

Species	n	%	
		Overall (n=400)	Among yeast colonized (n=360)
<i>C. tropicalis</i>	164	41.00	45.56
<i>C. albicans</i>	151	37.75	41.94
<i>C. parapsilosis</i>	23	5.75	6.39
<i>C. lusitaniae</i>	4	1.00	1.11
<i>C. glabrata</i>	5	1.25	1.39
<i>C. dubliniensis</i>	4	1.00	1.11
<i>C. famata</i>	3	0.75	0.83
<i>C. ciferrii</i>	2	0.50	0.56
<i>C. krusei</i>	2	0.50	0.56
<i>C. tropicalis</i> + <i>C. duobushaemulonii</i>	1	0.25	0.28
<i>C. parapsilosis</i> + <i>K. ohmeri</i>	1	0.25	0.28

Table 4. Distribution of the isolated mold species

Species	n	%	
		Overall (n=400)	Among mold colonized (n=28)
<i>A. fumigatus</i>	11	2.75	39.29
<i>A. oryzae</i>	6	1.50	21.43
<i>A. flavus</i>	6	1.50	21.43
<i>A. chevalieri</i>	2	0.50	7.14
<i>A. nomius</i>	2	0.50	7.14
<i>F. solani</i>	1	0.25	3.57

Table 5. Distribution of the species causing invasive fungal infection

Species	Fungal wound infection (n=39)		Fungemia (n=15)	
	n	%	n	%
<i>C. tropicalis</i>	13	33.33	10	66.67
<i>C. albicans</i>	7	17.95	3	20.00
<i>C. parapsilosis</i>	1	2.56	2	13.33
<i>K. ohmeri</i>	1	2.56		
<i>A. fumigatus</i>	4	10.26		
<i>A. flavus</i>	4	10.26		
<i>A. oryzae</i>	2	5.13		
<i>A. chevalieri</i>	1	2.56		
<i>A. nomius</i>	1	2.56		
<i>F. solium</i>	1	2.56		
<i>C. tropicalis</i> and <i>C. duobushaemulonis</i>	1	2.56		
<i>C. tropicalis</i> and <i>A. fumigatus</i>	1	2.56		
<i>C. tropicalis</i> and <i>A. flavus</i>	2	5.13		

There were 28 patients colonized or infected with molds, with *Aspergillus* species as the predominant agent (27/28), followed by *Fusarium*. The most common *Aspergillus* species was *A. fumigatus* (39.29%; Table 4).

The frequency of fungal species responsible for IFI is listed in Table 5. The most common agents of FWI were *Candida* (including *K. ohmeri*; n=26; 66.7% of FWI patients) and *Aspergillus* (n=15, 38.5%) species. *Candida tropicalis* was the most frequently isolated pathogen from the wound (n=17; 43.6% of FWI patients), followed by *C. albicans* (n=7; 17.95% of FWI patients) and *A. flavus* (n=6; 15.4% of FWI patients). Four patients were simultaneously overwhelmed by two different fungal species. Among three species of *Candida* causing candidaemia, *C. tropicalis* was the most prevalent species that accounted for 66.67% of the 15 yeast isolates. The other species were *C. albicans* (20.0%) and *C. parapsilosis* (14.33%).

The univariate analysis showed that severe infection, prolonged ICU stay, hyperglycemia, dialysis, and parenteral nutrition were significantly correlated with FC. However, in multivariate analysis, no factor was found to expose the patients to a higher risk of FC (Table 6). However, this analysis revealed many factors related to IFI. The results revealed hyperglycemia, prolonged ICU stay, and heavy *Candida* species colonization as the independent predictors of IFI (Table 7). Figure 1 shows the positive results of the specimens cultured within weekly intervals postburn. The frequency of IFI steadily increased during the stay in the hospital, while the prevalence of FC was nearly unchanged during the study period.

Table 6. Factors associated with fungal colonization

Variable	Univariate logistic regression		Multiple logistic regression	
	OR (95%CI)	P	OR (95%CI)	P
Severe injury	2.333 (1.207-4.512)	0.013	1.339 (0.629-2.852)	0.449
Inhalation injury	2.769 (0.829-9.252)	0.120		
Severe infection	2.407 (1.175-4.933)	0.028	1.620 (0.701-3.746)	0.259
Apache>20	3.545 (0.470-26.722)	0.088		
Prolonged ICU stay	1.512(0.781-2.926)	0.245		
Renal failure	2.449 (0.846-7.091)	0.061		
Hyperglycemia	2.818 (1.304-6.090)	0.004	1.355 (0.539-3.406)	0.518
Catheter	1.535 (0.785-3.004)	0.213		
Hemodialysis	7.645 (1.030-56.738)	0.018	4.758 (0.594-38.139)	0.142
Total parenteral nutrition	2.467 (1.258-4.836)	0.011	1.402 (0.626-3.138)	0.412
Mechanical ventilation	1.902 (0.850-4.256)	0.077		
Immunopressive therapy	1.753 (0.901-3.411)	0.099		

Table 7. Factors related to invasive fungal infection

Variable	Univariate logistic regression		Multiple logistic regression	
	OR (95%CI)	P	OR (95%CI)	P
Inhalation injury	1.576 (0.778-3.195)	0.233		
Severe injury	13.843 (3.311-57.885)	0.001	3.209 (0.626-16.458)	0.162
Burn shock	1.648 (0.376-7.229)	0.753		
Severe infection	12.996 (1.765-95.683)	0.001	0.255 (0.036-1.820)	0.173
Apache>20	1.350 (0.494-3.692)	0.573		
Prolonged ICU stay	4.520 (2.195-9.309)	0.001	2.611 (1.114-6.117)	0.027
Renal failure	3.729 (2.004-6.937)	0.001	1.101 (0.439-2.764)	0.837
Hyperglycemia	13.274 (5.509-31.982)	0.001	3.984 (1.309-12.129)	0.015
Catheter	8.491 (2.591-27.829)	0.001	0.951 (0.216-4.192)	0.947
Hemodialysis	6.723 (3.523-12.830)	0.001	2.071 (0.827-5.187)	0.120
TPN	19.800 (2.698-145.298)	0.001	4.724 (0.398-56.141)	0.219
Mechanical ventilation	3.875 (2.115-7.102)	0.001	0.696 (0.286-1.693)	0.424
Immunopressive therapy	8.364 (3.478-20.111)	0.001	2.418 (0.838-6.971)	0.102
Fungal colonization		0.005		0.997
Heavy fungal colonization	5.787 (2.406-13.920)	0.001	2.853 (1.095-7.437)	0.032

TPN: total parenteral nutrition




Figure 1. Prevalence of fungal colonization and infection in different weeks

Discussion

This study was targeted toward examining the prevalence and related factors of fungal colonization/infection to provide guidance on the prevention and treatment practices of fungal infection for ICU burn patients. To the best of our knowledge, this is the first study that assessed the risk factors for fungal colonization/infection in burn patients residing in ICU in Vietnam. Our study revealed that ICU burn patients have some risk factors for IFI, which seemed to be remarkable for this set of vulnerable patients.

In the present study, the diagnosis of fungal colonization or infection was based on the detection of fungi in clinical samples by microscopic examination or culture. Microscopic potassium hydroxide preparation was used as it facilitates the visualization of fungal elements during direct microscopic examination. The H&E staining is the most common type of histopathological examination. Furthermore, PAS staining is useful for screening fungi in tissue sections as it binds with carbohydrate-rich macromolecules commonly found in fungal cell walls. Culture remains one of the key methods for diagnosing fungal infection, and in case of suspected fungemia, blood culture is currently considered the “gold standard” [22]. The other benefit of culture is the identification of the causative agents by conventional or molecular methods. The definite diagnosis of IFI, in the present study, was provided with histological and cultural evidence from biopsies or blood as required [23].

The results showed that males (77.75%) were more affected by burn injury, compared to females; moreover, the most common age group involved was 1-10 years. These findings are in concordance with previous reports in Vietnam [24], as well as China, a developing, neighbor country [25, 26]. The mortality rate in the study was low (19.5%) and comparable to the value reported by Mundhada SG et al. (18%) [27], indicating the reducing trend of burn-related mortality in various regions [28].

Our findings showed that FC among the ICU burn patients was very common (90%), with 100% of the

colonized patients having *Candida* isolation. The high rate of *Candida* colonization among those in ICU is in accordance with the data obtained by Caggiano et al. (2011) reporting the colonization of *Candida* species in 92.3% of the patients 15 days after ICU admission [29]. The very high rates of colonization by *Candida* species expose the patients to a high risk of IFI because the responsible agents for invasive candidiasis were mainly endogenous in origin [7], [30].

The FWI rate obtained in the present study is comparable to those reported by Capoor et al. (10%) [5], Gupta et al. (10%) [7], and Ali et al. (13%) [31]. In the current research, The prevalence of fungemia was 3.75%, which is in line with previous results [32-34]. This finding is indicative of the higher risk of fungemia in ICU burn patients, compared to that in the patients residing in other types of ICUs [35, 36]. There is also a study reporting a higher incidence of fungemia in ICU burn patients (11%) [37].

In the present study, all colonized patients were found to be colonized with yeast species. This is comparable to previous reports describing the frequency of yeast (mostly *Candida* species) in burn patients [38-40]. It should be noted that the isolation rates of species belonging to CTG clade (*C. glabrata* and *C. krusei*) that were usually resistant to fluconazole were low (1.5% and 0.5%, respectively). This finding is in accordance with a previous study performed on Vietnamese patients [41]. In the present study, the results also revealed some other rare yeasts, such as *C. famata*, *C. duobushaemulonii*, and *Pichia (Kodamaea) ohmeri*. Those species have gained importance nowadays due to the emergence of nosocomial pathogens and their rapidly increased resistance to antifungal agents [42-46]. In line with the previous studies [40, 47], the rate of colonization caused by mold was low (7%).

Candida tropicalis was the predominant agent responsible for both FWI and fungemia. In Vietnam, there are no prior data regarding the prevalence of IFI in ICU patients for comparative purpose. However, our results are in line with those of previous research with *C. tropicalis* being the most common agent causing

candidaemia (50.5%) [41]. In concordance with the data reported in an Iranian study [31], in the present research, more than 40% of FWIs were caused by molds. The high rate of FWI caused by mold is disproportionate to the low prevalence of mold colonization among burn patients (7%). This signifies the importance of measures to protect the wound from mold colonization [48]. The source of mold infection is mostly exogenous when conidia as filamentous fungi fall onto the wound from the surrounding air and invade the wound immediately [8]. The most common mold was *Aspergillus* species (15/16 patients). This may demonstrate the strong virulence of *Aspergillus* among many other molds [49]. It is worth noting that there was an FWI case infected with *F. solani*. Although invasive infection caused by *Fusarium* is rare, it has caught much attention for its angioinvasive property, resistance to conventional antifungal therapy [50], and high mortality rate [51].

Consistent with other reports [29, 52, 53], the univariate analysis revealed severe infection, prolonged ICU stay, hyperglycemia, dialysis, and parenteral nutrition as the significant predictors of FC. However, based on the multivariate analysis, there were no factors predicting FC in burn patients. In our opinion, this is reasonable because most of the colonized patients had isolation upon the ICU admission, and the rate of FC was nearly steady over the period of the patients' ICU stay (Figure 1). Hedderwick et al. (2000) also found that most of the colonized patients were already colonized with yeast upon admission to the ICU, and only some of them became colonized after admission [54].

The results of our study showed that hyperglycemia, prolonged ICU stay, and heavy *Candida* species colonization were independently predictive of IFI. Hyperglycemia during acute hospitalization is a common pathophysiological phenomenon among severely burned patients [55]. Hyperglycemia is an adaptive response but could lead to some adverse outcomes, especially with increased risk of infections, including fungal infection [5, 56-61]. In the current study, hyperglycemia was observed in 42.75% of the patients. Moreover, those with hyperglycemia were at 3.984 times higher (adjusted) risk of IFI, compared with those without hyperglycemia. Multiple site colonization is a well-known risk factor for invasive fungal infection and has been mentioned in many established *Candida* Prediction Rules [57, 62-65]. In line with the results of previous studies [53, 66, 67], prolonged ICU stay was also found to be an important factor; in this regard, there was a clear rise in the infection rate from the second week (Figure 1) of hospital stay.

The other factors found to be related to IFI in univariate analysis included severe injury, severe infection, renal failure, hemodialysis, total parenteral nutrition, mechanical ventilation, catheter, and immunosuppressive therapy. However, these factors were not significant in multivariate analysis.

Although these factors present in some *Candida* Prediction Rules, their impact may not be significant as they are absent in some other rules [57, 62-65]. Therefore, their contribution as confounding factors in this study was rational. In the present research, high APACHE II score was not associated with the increased risk of IFI, which is in line with the findings reported by Escrig AIR et al. (2016) [68]. These findings are significant for clinicians to take timely precaution and prophylactic practices against fungal infections for high-risk patients.

Conclusion

In conclusion, severely burned patients are at a risk of fungal infection mainly because of the pathophysiological consequences of injury and longer hospital stay. The most common agent was found to be *Candida* species (especially *C. tropicalis*), followed by *Aspergillus*. To ensure early and appropriate management measures, it is required to perform close monitoring on the presence of microbial agents in burn patients. It is also needed to well control blood glucose concentration in burn patients, especially in those staying in ICU for a long period.

Acknowledgments

The authors are grateful to Ms. Linh Khanh Le for revising the English text.

Author's contribution

N. V. B., N. T. X., and L. T. A contributed to the conception and design of the study, interpretation of the data, and final editing of the manuscript. D. X. Q. and C. B. L. contributed to data analysis and preparation of the manuscript draft. N. T. N. M. and N. N. L collected the clinical data and preparation of clinical samples. D. N. A., T. T. T. H and H. X. S involved in laboratory work. All of the authors agree to submit the manuscript for possible publication in the "Current Medical Mycology".

Conflicts of interest

The authors declare no conflicts of interest regarding the publication of this paper.

Financial disclosure

This work was partially supported by the Vietnam Ministry of National Defence (grant No. 2017.75.054, to TAL). The funding agency does not have any role in data collection, analysis, and writing of the manuscript. This research is part of a thesis submitted in partial fulfilment of the requirement for a Ph.D. degree in Literature and Philosophy in Health at the National Institute of Malaria, Parasitology, and Entomology of Vietnam.

References

1. World Health Organization. Burns. Geneva: World Health Organization; 2018.
2. Church D, Elsayed S, Reid O, Winston B, Lindsay R. Burn wound infections. Clin Microbiol Rev. 2006; 19(2):403-34.

3. Cochran A, Morris S, Edelman L, Saffle J. Systemic *Candida* infection in burn patients: a case-control study of management patterns and outcomes. *Surg Infect*. 2002; 3(4):367-74.
4. Struck MF, Gille J. Fungal infections in burns: a comprehensive review. *Ann Burns Fire Disasters*. 2013; 26(3):147-53.
5. Capoor MR, Sarabahi S, Tiwari VK, Narayanan RP. Fungal infections in burns: diagnosis and management. *Indian J Plast Surg*. 2010; 43(Suppl):S37-42.
6. Lachiewicz AM, Hauck CG, Weber DJ, Cairns BA, van Duin D. Bacterial infections after burn injuries: impact of multidrug resistance. *Clin Infect Dis*. 2017; 65(12):2130-6.
7. Gupta N, Haque A, Lattif AA, Narayan RP, Mukhopadhyay G, Prasad R. Epidemiology and molecular typing of *Candida* isolates from burn patients. *Mycopathologia*. 2004; 158(4):397-405.
8. Mousa H, Al-Bader S, Hassan D. Correlation between fungi isolated from burn wound and burn care units. *Burns*. 1999; 25(2):145-7.
9. Garcia-Hermoso D, Criscuolo A, Lee SC, Legrand M, Chaouat M, Denis B, et al. Outbreak of invasive wound mucormycosis in a burn unit due to multiple strains of *Mucor circinelloides* f. *circinelloides* resolved by whole-genome sequencing. *mBio*. 2018; 9(2):e00573-18.
10. Cornely O, Alastruey-Izquierdo A, Arenz D, Chen SC, Dannaoui E, Hochhegger B, et al. Global guideline for the diagnosis and management of mucormycosis: an initiative of the European Confederation of Medical Mycology in cooperation with the Mycoses Study Group Education and Research Consortium. *Lancet Infect Dis*. 2019; 19(12):e405-21.
11. Sarabahi S, Tiwari VK, Arora S, Capoor MR, Pandey A. Changing pattern of fungal infection in burn patients. *Burns*. 2011; 38(4):520-8.
12. Atiyeh B, Masellis A, Conte C. Optimizing burn treatment in developing low- and middle-income countries with limited health care resources (Part 1). *Ann Burn Fire Disasters*. 2009; 22(3):121-5.
13. Wallace AB. The exposure treatment of burns. *Lancet*. 1951; 1(6653):501-4.
14. Knaus W, Zimmerman J, Wagner D, Draper EA, Lawrence DE. APACHE-acute physiology and chronic health evaluation: a physiologically based classification system. *Crit Care Med*. 1981; 9(8):591-7.
15. Greenhalgh DG, Saffle JR, Holmes JH 4th, Gamelli RL, Palmieri TL, Horton JW, et al. American burn association consensus conference to define sepsis and infection in burns. *J Burn Care Res*. 2007; 28(6):776-90.
16. Leaw SN, Chang HC, Sun HF, Barton R, Bouchara JP, Chang TC. Identification of medically important yeast species by sequence analysis of the internal transcribed spacer regions. *J Clin Microbiol*. 2006; 44(3):693-9.
17. Horvath E, Murray C, Vaughan G, Chung KK, Hospenthal DR, Wade CE, et al. Fungal wound infection (not colonization) is independently associated with mortality in burn patients. *Ann Surg*. 2007; 245(6):978-85.
18. Pittet D, Monod M, Suter PM, Frenk E, Auckenthaler R. *Candida* colonization and subsequent infections in critically ill surgical patients. *Ann Surg*. 1994; 220(6):751-8.
19. Bellomo R, Ronco C, Kellum JA, Mehta RL, Palevsky P; Acute dialysis quality initiative workgroup. acute renal failure - definition, outcome measures, animal models, fluid therapy and information technology needs: the Second International Consensus Conference of the Acute Dialysis Quality Initiative (ADQI) Group. *Crit Care*. 2004; 8(4):R204-12.
20. Rhodes A, Evans L, Alhazzani W, Levy MM, Antonelli M, Ferrer R, et al. Surviving sepsis campaign: international guidelines for management of sepsis and septic shock: 2016. *Intensive Care Med*. 2017; 43(3):304-77.
21. Arabi Y, Venkatesh S, Haddad S, Al Shimemeri A, Al Malik S. A prospective study of prolonged stay in the intensive care unit: predictors and impact on resource utilization. *Int J Qual Heal Care*. 2002; 14(5):403-10.
22. Ostrosky-zeichner L. Invasive mycoses: diagnostic challenges. *Am J Med*. 2012; 125(1 Suppl):S14-24.
23. De Pauw B, Walsh TJ, Donnelly JP, Stevens DA, Edwards JE, Calandra T, et al. Revised definitions of invasive fungal disease from the European Organization for Research and Treatment of Cancer/ Invasive Fungal Infections Cooperative Group and the National Institute of Allergy and Infectious Diseases Mycoses Study Group (EORTC/MSG). *Clin Infect Dis*. 2008; 46(12):1813-21.
24. Lam NN, Huong HT, Tuan CA. Mass burn injuries: an analysis of characteristics and outcomes in a developing country. *Ann Burn Fire Disasters*. 2017; 30(3):210-3.
25. Tian H, Wang L, Xie W, Shen C, Guo G, Liu J, et al. Epidemiologic and clinical characteristics of severe burn patients: results of a retrospective multicenter study in China, 2011-2015. *Burn Trauma*. 2018; 6:14.
26. Li H, Yao Z, Tan J, Zhou J, Li Y, Wu J, Luo G. Epidemiology and outcome analysis of 6325 burn patients: a five-year retrospective study in a major burn center in Southwest China. *Sci Rep*. 2017; 7:46066.
27. Mundhada SG, Waghmare PH, Rathod PG, Ingole KV. Bacterial and fungal profile of burn wound infections in Tertiary Care Center. *Indian J Burns*. 2015; 23(1):71-5.
28. Smolle C, Cambiaso-Daniel J, Forbes AA, Wurzer P, Hundeshagen G, Branski LK, et al. Recent trends in burn epidemiology worldwide: a systematic review. *Burns*. 2017; 43(2):249-57.
29. Caggiano G, Puntillo F, Coretti C, Giglio M. *Candida* colonization index in patients admitted to an ICU. *Int J Mol Sci*. 2011; 12(10):7038-47.
30. Cole GT, Halawa AA, Anaissie EJ. The role of the gastrointestinal tract in hematogenous candidiasis: from the laboratory to the bedside. *Clin Infect Dis*. 1996; 22(Suppl 2):S73-88.
31. Bahar MA, Pakyari MR, Gholipourmalekabadi M, Samadikuchaksaraei A. The prevalence of fungal infections in a level I Iranian burn hospital. *Asian Biomed*. 2013; 7(6):829-33.
32. Spebar M, Pruitt BJ. Candidiasis in the burned patient. *J Trauma*. 1981; 21(3):237-9.
33. Ha JF, Italiano CM, Heath CH, Shih S, Rea S, Wood FM. Candidemia and invasive candidiasis: A review of the literature for the burns surgeon. *Burns*. 2011; 37(2):181-95.
34. Vinsonneau C, Benyamina M, Baixench MT, Stephanazzi J, Augris C, Grabar S, et al. Effects of candidaemia on outcome of burns. *Burns*. 2009; 35(4):561-4.
35. Bassetti M, Giacobbe DR, Vena A, Trucchi C, Ansaldi F, Antonelli M, et al. Incidence and outcome of invasive candidiasis in intensive care units (ICUs) in Europe: results of the EUCANDICU project. *Crit Care*. 2019; 23(1):219.
36. Ghrenassia E, Mokart D, Mayaux J, Demoule A, Rezine I, Kerhuel L, et al. Candidemia in critically ill immunocompromised patients: report of a retrospective multicenter cohort study. *Ann Intensive Care*. 2019; 9(1):62.
37. Fochtmann A, Forstner C, Haggmann M, Keck M, Muschitz G, Presterl E, et al. Predisposing factors for candidemia in patients with major burns. *Burns*. 2015; 41(2):326-32.
38. Fitzwater J, Purdue GF, Hunt JL, O'Keefe GE. The risk factors and time course of sepsis and organ dysfunction after burn trauma. *J Trauma Acute Care Surg*. 2003; 54(5):959-66.
39. Ballard JL, Edelman L, Saffle J, Sheridan R, Kagan R, Bracco D, et al. Positive fungal cultures in burn patients: a multicenter review. *J Burn Care Res*. 2008; 29(1):213-21.
40. Rafik A, Diouri M, Bahechar N, Chlihi A. Epidemiology of nosocomial fungal infections in the National center for burns in Casablanca, Morocco. *Ann Burns Fire Disasters*. 2016; 29(2):90-3.
41. Bac ND, Anh LT, Quang LB, Luc NK, Nga TT, Nagi M, et al. Prevalence of *Candida* bloodstream isolates from patients in two hospitals in Vietnam. *Iran J Microbiol*. 2019; 11(2):108-13.
42. Beyda ND, Chuang SH, Alam MJ, Shah ND, Ng TM, McCaskey L, et al. Treatment of *Candida* famata bloodstream infections: case series and review of the literature. *J Antimicrob Chemother*. 2013; 68(2):438-43.
43. Boatto HF, Cavalcanti SD, Del Negro GM, Girão MJ, Francisco EC, Ishida K, et al. *Candida* duobushaemulonii: an emerging rare pathogenic yeast isolated from recurrent vulvovaginal candidiasis in Brazil. *Mem Inst Oswaldo Cruz*. 2016; 111(6):407-10.
44. Ramos R, Caceres D, Perez M, Garcia N, Castillo W, Santiago

- E, et al. Emerging multidrug-resistant *Candida duobushaemulonii* infections in Panama hospitals: importance of laboratory surveillance and accurate identification. *J Clin Microbiol.* 2018; 56(7):e00371.
45. Hou X, Xiao M, Chen S, Wang H, Cheng JW, Chen XX, et al. Identification and antifungal susceptibility profiles of *Candida haemulonii* species complex clinical isolates from a multicenter study in China. *J Clin Microbiol.* 2016; 54(11):2676-80.
 46. Chakrabarti A, Rudramurthy S, Kale P, Hariprasath P, Dhaliwal M, Singhi S, et al. Epidemiological study of a large cluster of fungaemia cases due to *Kodamaea ohmeri* in an Indian tertiary care centre. *Clin Microbiol Infect.* 2014; 20(2):O83-9.
 47. Ibrahim N, Amer T. Frequency of bacterial and fungal infections of burn wounds at cairo university burn center. *Egypt J Med Microbiol.* 2008; 17(4):573-83.
 48. Guinea J, de Viedma DG, Peláez T, Escribano P, Muñoz P, Meis JF, et al. Molecular epidemiology of *Aspergillus fumigatus*: an in-depth genotypic analysis of isolates involved in an outbreak of invasive *Aspergillosis* downloaded from <http://jcm.asm.org/> on September 30, 2018 by guest. *J Clin Microbiol.* 2011; 49(10):3498-503.
 49. Enoch DA, Yang H, Aliyu SH, Micallef C. The changing epidemiology of invasive fungal infections. *Methods Mol Biol.* 2017; 1508:17-65.
 50. Muhammed M, Coleman J, Carneiro H, Mylonakis E. The challenge of managing fusariosis. *Virulence.* 2011; 2(2):91-6.
 51. Lass-Flörl C, Cuenca-Estrella M. Changes in the epidemiological landscape of invasive mold infections and disease. *J Antimicrob Chemother.* 2017; 72(Suppl 1):i5-11.
 52. Goyal NK, Gore MA, Goyal RS. Fungal colonisation in burn wounds: an Indian scenario. *Indian J Surg.* 2010; 72(1):49-52.
 53. Sharma S, Bajaj D, Sharma P. Fungal infection in thermal burns: a prospective study in a tertiary care centre. *J Clin Diagn Res.* 2016; 10(9):PC05-7.
 54. Hedderwick SA, Lyons MJ, Liu M, Vazquez JA, Kauffman CA. Epidemiology of yeast colonization in the intensive care unit. *Eur J Clin Microbiol Infect Dis.* 2000; 19(9):663-70.
 55. Jeschke MG, Chinkes DL, Finnerty CC, Kulp G, Suman OE, Norbury WB, et al. Pathophysiologic response to severe burn injury. *Ann Surg.* 2008; 248(3):387-401.
 56. Hemmila MR, Taddonio MA, Arbabi S, Maggio PM, Wahl WL. Intensive insulin therapy is associated with reduced infectious complications in burn patients. *Surgery.* 2008; 144(4):629-37.
 57. Paramythiotou E, Frantzeskaki F, Flevari A, Armaganidis A. Invasive fungal infections in the ICU: how to approach, how to treat. *Molecules.* 2014; 19(1):1085-119.
 58. Shoham S, Han G, Granek T, Walsh T, Magee MF. Association between blood glucose levels and development of candidemia in hospitalized patients. *Endocr Pract.* 2009; 15(2):111-5.
 59. Sydnor ERM, Perl TM. Hospital epidemiology and infection control in acute-care settings. *Clin Microbiol Rev.* 2011; 24(1):141-73.
 60. Karacaer Z, Oncul O, Turhan V, Gorenk L, Ozyurt M. A surveillance of nosocomial candida infections: epidemiology and influences on mortality in intensive care units. *Pan Afr Med J.* 2014; 19:398.
 61. Prasad R. *Candida albicans*: cellular and molecular biology. Berlin, Germany: Springer; 2017.
 62. Rüping MJ, Vehreschild JJ, Cornely DO. Patients at high risk of invasive fungal infections: when and how to treat. *Drugs.* 2008; 68(14):1941-58.
 63. Eggimann P, Ostrosky-Zeichner L. Early antifungal intervention strategies in ICU patients. *Curr Opin Crit Care.* 2010; 16(5):465-9.
 64. Perlroth J, Choi B, Spellberg B. Nosocomial fungal infections: epidemiology, diagnosis, and treatment. *Med Mycol.* 2007; 45(4):321-46.
 65. Yapar N. Epidemiology and risk factors for invasive candidiasis. *Ther Clin Risk Manag.* 2014; 10:95-105.
 66. Raz-Pasteur A, Hussein K, Finkelstein R, Ullmann Y, Egozi D. Blood stream infections (BSI) in severe burn patients—early and late BSI: a 9-year study. *Burns.* 2013; 39(4):636-42.
 67. Van Burik JA, Colven R, Spach DH. Cutaneous aspergillosis. *J Clin Microbiol.* 1998; 36(11):3115-21.
 68. Renau Escrig AI, Salavert M, Vivó C, Cantón E, Pérez del Caz MD, et al. Candidemia in major burns patients. *Mycoses.* 2016; 59(6):391-8.